



King's Research Portal

DOI:

[10.1016/j.ceb.2015.12.001](https://doi.org/10.1016/j.ceb.2015.12.001)

Document Version

Publisher's PDF, also known as Version of record

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Olmos, Y., & Carlton, J. G. (2016). The ESCRT machinery: new roles at new holes. *Current Opinion in Cell Biology*, 38, 1–11. [10.1016/j.ceb.2015.12.001](https://doi.org/10.1016/j.ceb.2015.12.001)

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

The ESCRT machinery: new roles at new holes

Y Olmos and JG Carlton

The ESCRT machinery drives a diverse collection of membrane remodeling events, including multivesicular body biogenesis, release of enveloped retroviruses and both reformation of the nuclear envelope and cytokinetic abscission during mitotic exit. These events share the requirement for a topologically equivalent membrane remodeling for their completion and the cells deployment of the ESCRT machinery in these different contexts highlights its functionality as a transposable membrane-fission machinery. Here, we will examine recent data describing ESCRT-III dependent membrane remodeling and explore new roles for the ESCRT-III complex at the nuclear envelope.

Address

Division of Cancer Studies, Section of Cell Biology and Imaging, King's College London, London SE1 1UL, United Kingdom

Corresponding author: Carlton, JG (Jeremy.Carlton@kcl.ac.uk)

Current Opinion in Cell Biology 2016, **38**:1–11

This review comes from a themed issue on **Cell architecture**

Edited by **Matthieu Piel** and **Margaret Gardel**

For a complete overview see the [Issue](#) and the [Editorial](#)

Available online 15th January 2016

<http://dx.doi.org/10.1016/j.ceb.2015.12.001>

0955-0674/© 2016 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

Introduction


The Endosomal Sorting Complex Required for Transport (ESCRT) machinery is an evolutionarily-conserved, multi-subunit membrane remodeling complex (Table 1). Originally identified in yeast for its essential role in the biogenesis of intraluminal vesicles (ILVs) upon a class of endosome called the multivesicular body (MVB) [1–3], its roles in mammalian cells have been expanded to encompass a number of topologically equivalent membrane remodeling events. ILV biogenesis requires the endosome's limiting membrane to be pushed away from the cytosol to generate a nascent bud connected via a membranous stalk to the endosomal limiting membrane. Subsequent severing of this nascent ILV from the limiting membrane requires a membrane fission activity that acts on the cytosolic face of this membranous stalk (Figure 1a). It is thought that the ESCRT-III complex provides this activity, whilst upstream ESCRT-components (ESCRT-I and ESCRT-II) coordinate this fission with the forming and sorting of ubiquitinated cargo onto the nascent ILV [4,5]. As well as being a prerequisite to lysosomal degradation, the contents of MVBs can be released upon fusion of the

MVB with the plasma membrane — the released ILVs are exosomes and as their formation is ESCRT-dependent, ESCRTs thus offer a layer of control over this route of intercellular communication [6,7]. Whilst an endosomal function for ESCRTs has been demonstrated across a range of organisms, the topologically unique membrane remodeling that ESCRT-III complex can perform has allowed this machinery to be co-opted by a number of cellular factors and to drive a diverse range of physiological and pathophysiological cellular processes across organisms from *Archaea* to animals [8].

In mammalian cells, ESCRT-components have been implicated as host-factors for the release of enveloped retroviruses such as HIV-1 or Ebola [9,10,11,12]. The structural proteins of these enveloped viruses encode peptide-ligands (Late, or L-domains), that are able to directly bind ESCRT-proteins. Here, the ESCRT-III complex, recruited via TSG101 and ESCRT-I or by the ESCRT-III accessory protein ALIX [13,14], functions to sever the membranous stalks connecting budded virions with the plasma membrane in a manner topologically analogous to the membrane fission reaction performed to release ILVs at the MVB (Figure 1a). Whilst largely implicated in the budding of virions from the plasma membrane, ESCRT proteins have also been involved in the lifecycle and release of viruses such as Epstein-Barr virus that traverse intracellular membranes such as the nuclear envelope [15] and Herpes Simplex viruses [16] that buds through the nuclear envelope and again into the lumen of membranes organelles during secondary envelopment, suggesting that ESCRT-III may again be recruited for the membrane remodeling allowing these envelopment events.

In addition to a pathophysiological role in viral replication, the ESCRT proteins TSG101 and ALIX are recruited to the midbody of dividing cells through interaction with the midbody protein CEP55 [17,18,19]. Here, they function to recruit the ESCRT-III complex to perform a topologically equivalent membrane fission during cytokinetic abscission (Figure 1a), allowing the separation of daughter cells and the completion of cell division [18,20,21]. ESCRT-III proteins have additionally been implicated in the repair of damaged regions of the plasma membrane, where they are recruited in a Ca^{++} -dependent manner by ALG2 [22,23] to sever protruding, damaged regions of the plasma membrane in a manner similar to that employed in viral release. Further, ESCRT-III has been shown to play roles in autophagy [24,25], neuronal pruning [26], material transfer at the immunological synapse [27] and release of exosomes [6]. This article will not focus on these

Figure 1



Sites and models of ESCRT-III activity. **(a)** Cartoon depicting sites of topologically-equivalent membrane remodeling performed by the ESCRT-III complex. In all cases, ESCRT-III provides an activity allowing resolution of the membranous stalk with the concomitant separation of the two membranes previously connected by the stalk. This separation achieves release of ILVs (1), enveloped viruses such as HIV-1 (2), daughter cells (3) and separation of previously connected inner and outer nuclear membranes (4). Bottom schematic depicts membrane separation in each case achieved through ESCRT-III activity. **(b)** Models for ESCRT-III driven membrane fission. Filaments of polymerised ESCRT-III subunits are thought to assemble inside a membranous stalk, connecting two parental membranes. ESCRT-III assembly, either via the shape of the formed holo-polymer (dome model), or through constriction of the ESCRT-III filament (purse string model), narrows the stalk. This narrowing presumably makes

Table 1

ESCRT subunits in yeast and mammals Tabular list of ESCRT components in yeast and mammalian cells; in mammalian cells, ESCRT-III components are referred to as CHMPs, due to their identification as *Charged Multivesicular Body Proteins* [32]/*Chromatin Modifying Proteins* (CHMPs) [33]

Complex	Yeast	Mammals
ESCRT-I	Vps23	TSG101
	Vps28	VPS28
	Vps37	VPS37A, VPS37B, VPS37C, VPS37D
	Mvb12	MVB12A, MVB12B, UBAP1
ESCRT-II	Vps22	EAP30
	Vps25	EAP20
	Vps36	EAP45
ESCRT-III	Did2/Vps46	CHMP1A, CHMP1B
	Vps2	CHMP2A, CHMP2B
	Vps24	CHMP3
	Snf7	CHMP4A, CHMP4B, CHMP4C
	Vps60	CHMP5
	Vps20	CHMP6
	Chm7	CHMP7
	Vps4	VPS4A, VPS4B
	Vta1	LIP5
ESCRT associated	Bro1	ALIX, HD-PTP
	Ist1	IST1
	?	SPASTIN

classical (or even those more esoteric) functions of the ESCRT-machinery; readers are referred to myriad excellent reviews on this subject elsewhere [28–31]. Rather, we will focus on newly published data describing new roles for ESCRT-III during cell division.

ESCRT-III

ESCRT-III is the hypothesised membrane fission complex within the ESCRT-machinery and is directed to its sites of action through interaction with upstream ESCRT-components, themselves localized by various adaptor proteins as described above. ESCRT-III subunits (CHMPs, for *Charged Multivesicular Body Proteins* [32], or *Chromatin Modifying Proteins* [33]) transition between soluble and polymerising states, and assemble in a defined order to form a membrane-remodeling filament that brings about membrane fission. CHMP proteins have long been proposed to exist in closed and open conformations, with the hypothesis that the open conformation represented a filament-forming state. Recent cryo-EM data has at last resolved the molecular structure of a CHMP (CHMP1B) in the ‘open’ conformation and

revealed it to adopt an extended α -helical V-shaped structure that bound to proximal open-forms of itself in a regular array to form a filament in a co-polymer with the N-terminal domain of IST1 [34**]. Interestingly, IST1 in this filament was found on the outside of this filament in the closed conformation, whereas CHMP1B in this filament was found in the open conformation. Of note, CHMP1B and IST1 play atypical roles in ESCRT-biology, being dispensable for HIV-1 release and degradative cargo sorting and whilst IST1 functions in cytokinesis [35,36] and at the nuclear envelope [37**] (see below), IST1 is also necessary for scission of recycling carriers leaving endosomes [38]. Consistent with this, the CHMP1B:IST1 filament contained a luminal (rather than peripheral) membrane-binding surface, suggesting a function as a scaffold around membranous stalks, rather than inside them [34**]. Indeed, the IST1:CHMP1B filaments were straight and not spiral indicating that they may be better suited to constriction of a tubule rather than the tightening of a stalk (as described below). These data indicate that a heteropolymeric ESCRT-III filament can accommodate CHMPs in different activation states and suggest that the open form represents the membrane-binding conformation. The structure of an ‘open’ CHMP will hopefully guide our understanding of how filaments formed from CHMP-2, CHMP-4 and CHMP-3 generate and stabilise the negative curvature required for ESCRT-III dependent membrane separation. Mechanisms of ESCRT-III assembly and disassembly have been reviewed well elsewhere [29,31,39] and models of how ESCRT-III achieves the terminal membrane fission have been proposed [39–41]. These models of ESCRT-III function remain largely untested, but are linked by the basic premise that ESCRT-III assembles as a spiral filament on the cytosolic face of a membranous stalk, and provides an activity allowing resolution of this stalk and separation of the two previously connected membranes (Figure 1b). Exactly how this scission occurs is not known, but remodeling of the ESCRT-III polymer may contribute either to membrane deformation, or to the membrane separation. It is known that ESCRT-III proteins form spiral filaments in vitro, suggesting that the oligomer may have a preferred curvature. It was recently shown that filaments formed from Vps32 (the *C. elegans* homologue of CHMP4B/Snf7) could accommodate a range of bending angles [42**] and analysis of Snf7 filament assembly on supported bilayers have added some new considerations into how filament assembly may drive membrane remodelling. Using a combination of high-speed atomic force microscopy and electron

(Figure 1 Legend Continued) it energetically more favourable to separate the membranes, rather than persist with membranes connected by a highly-curved, thin, membranous stalk. Models propose that the AAA-ATPase VPS4 acts either to tighten the filament, through sequential extraction of polymerised CHMPs (purse string model, VPS4 activity needed throughout remodeling event (1)) or through the induction of conformational changes in the CHMPs by direct interaction. Alternatively, membrane fission is accomplished through formation of the ESCRT-III holo-polymer (dome model, with VPS4 acting after fission to disassemble ESCRT-III filaments for subsequent rounds of assembly (2)). Period of VPS4 activity indicated by arrow.

microscopy, circular arrays of Snf7 with a preferred diameter of 25 nm were observed to break and seed polymerisation of helical Snf7 filaments that grew radially on supported lipid membranes [43^{••}]. As these spirals grew, the degree of filament curvature was necessarily reduced to accommodate the growing spiral, putting elastic strain on the filament. The growing Snf7 filament was suggested to act as a spiral-spring, storing elastic energy as the filament extended; indeed, molecular dynamics simulations predict the free energy state of Vps32 filaments is dramatically increased at the edge of the spiral [42^{••}] and measurements of membrane tension [43^{••}] were increased upon Snf7 polymerisation. Upon reaching confluency, spiral growth was inhibited by lateral interactions, and spirals released their stored energy through deformation, suggesting that in a cellular environment, this release of elastic energy could be transferred to the membrane. The authors propose that this energy drives ESCRT-III-dependent membrane deformation (Figure 2), perhaps by moving leaflets of the bilayer relative to each other. Whilst this is a likely scenario, it should also be noted that ESCRT-III functions in a number of situations where membrane deformation is dispensable (for example, during virion release, nuclear envelope sealing and cytokinetic abscission; here the membranous stalk to be resolved is pre-existing) and it is possible that the energy stored in these filaments is alternatively used to drive membrane separation rather than membrane deformation. An additional point that these data raise is a consideration of scale. Whilst viral stalks, ILV stalks, and stalks to be resolved in the nuclear envelope are of similar dimensions (30–50 nm), the midbody is much larger, with ESCRT-III components being


observed when the midbody has diameters of approximately 1 μm (Figure 3b [44]). This will necessitate the formation of a filament with a very low curvature, that would be predicted to store great deal of elastic energy, perhaps (given the larger diameter of ESCRT-III-dependent cytokinetic filaments [44]) necessitating partner proteins to prevent breakage, and be capable of transferring this greater energy to the membrane which may be necessary for efficient constriction and abscission of this larger structure.

In addition to individual ESCRT-III subunits, the MIT-domain containing AAA-ATPase, VPS4, is recruited to short peptide motifs in the C-terminus of ESCRT-III subunits to allow for ESCRT-III filament disassembly and subunit reuse in subsequent rounds of ESCRT-activity — VPS4 binding could also remodel and constrict the Vps32 spirals in an ATPase-independent manner [42^{••}], suggesting that it too could modulate the elastic energy within the ESCRT-III filament. Interestingly, ESCRT-III subunits are additionally capable of recruiting alternate AAA-ATPases, such as the microtubule-severing enzyme Spastin [45], which have important roles in coordinating ESCRT-III-dependent membrane and cytoskeletal modeling as described below.

ESCRTs on the nuclear envelope — NPC surveillance


The Nuclear Pore Complex (NPC) is an ancient, conserved and long-lived structure that allows gated exchange between nucleoplasm and cytoplasm, allowing the establishment of proper nucleo-cytoplasmic compartmentalization [46]. A recent epistasis screen uncovered an unexpected role for the core ESCRT-III complex in extracting defective NPCs, thus ascribing a surveillance role for this complex at the nuclear envelope [47^{••}]. In this role, ESCRT-III was recruited to the nuclear envelope through interaction of Snf7 with the *Lap2-Emerin-MAN1* (LEM) family proteins Heh1 and Heh2 that had previously been implicated in NPC quality control [48]. In yeast lacking ESCRT-III components, Vps4 or Heh2, aberrantly assembled NPC components were sequestered within a domain on the nuclear envelope termed the SINC (Storage of Improperly assembled NPCs). Unlike mammalian cells who disassemble and reassemble their nuclear envelope during each open mitosis, yeast undergo a closed mitosis and thus require surveillance mechanisms such as ESCRT-III (and parallel quality-control mechanisms such as the ASI-complex may tie the ERAD pathway into surveillance of other ubiquitinated proteins at this organelle [49,50]), to extract malfunctioning NPC components. Whether a similar surveillance role for ESCRT-III exists in higher organisms remains to be established, but is likely to be complicated by the open-mitosis that occur in these systems. The basis for ESCRT-III dependent extraction of defective Nups likely involves ubiquitin as proteosomal (but not vacuolar) inhibition impaired this extraction [47^{••}],

Figure 2



Energetic considerations in ESCRT-III assembly. Cartoon depicting the energy stored in helical polymers of ESCRT-III subunits and the suggestion that this energy is released to allow membrane deformation.

Figure 3



Mitotic ESCRT-III localization. Immunofluorescence analysis of HeLa cells stained with antibodies against endogenous CHMP2A or tubulin as indicated, demonstrating CHMP2A localization to the reforming nuclear envelope (a) or the midbody during cytokinesis (b). For methods and more detailed images, see Olmos *et al.* (2015). In A, dotted outline depicts cell border, images obtained by deconvolution of widefield images (a) or confocal imaging (b). Scale bar is 10 μ m.

however, the exact mechanism by which ESCRT-III extracts membrane proteins and directs them to the proteasome (rather than the lysosome) is unknown. Whilst the ESCRT-machinery commonly processes ubiquitinated proteins, the absence of ubiquitin-recognition domains within ESCRT-III suggests that additional components of this extraction machinery that direct ubiquitinated nucleoporins for ESCRT-III mediated extraction may exist. Webster and Lusk have proposed attractive models for the operation of ESCRT-III in this context [51] — given the role of AAA-ATPases such as VPS4 in disassembling and solubilizing protein complexes, it is possible that recruited Vps4 plays a direct role in the extraction of these defective intermediates, or uses the physical link between ESCRT-III bound nucleoporins to extract them indirectly,

via extraction of the ESCRT-III subunit. Alternatively, budding profiles at the INM and intra-membrane vesicular structures were observed in $\delta vps4$ cells suggesting that these defective nucleoporins may be cleared through vesicular-budding into the intermembrane space [47^{••}]. Whilst this model is topologically satisfying from an ESCRT-III perspective, it is unclear how vesicles in the inter-membrane space could then access the proteasome for degradation.

ESCRT-III in nuclear envelope reformation

Cell division in eukaryotes involves extensive remodeling of the nuclear envelope (NE) to ensure proper segregation of nuclear and cytoplasmic contents. Higher eukaryotic cells divide by open mitosis, in which the nuclear envelope


is broken down at mitotic onset to allow the chromosomes to access the mitotic spindle. This process involves the disassembly and dispersal of all the main elements of the nuclear envelope, including the nuclear membranes, NPCs and the lamina. Upon mitotic onset, some ESCRT-III subunits are subject to phosphorylation, which may render them inactive until they are dephosphorylated during mitotic exit [52]. After chromosome segregation, during telophase, a new nuclear envelope is reformed around each daughter nuclei in order to re-establish nucleocytoplasmic partitioning [53–56]. Nuclear envelope reformation is a two-step process that requires extensive membrane fusion. First, ER membranes, proposed to be either cisternal [57,58] or tubular [59,60], attach to chromosomes and spread out, coating the chromatin surface in double-membrane sheets. Secondly, and in order to form a completely sealed nuclear envelope, it is necessary to close any remaining holes or gaps within the double nuclear membranes through the process of annular fusion [55]. In vitro studies of nuclear envelope reassembly in *Xenopus* egg extracts showed that both steps of nuclear envelope reformation are regulated by the p97 AAA-ATPase in association with its adaptor proteins p47, in the case of membrane delivery and nuclear expansion, or UFD1/NPL4 for the annular fusion event [61]. The mechanism by which annular fusion occurs has remained largely unknown, but the topology of this process is identical to cytokinetic abscission as the closure of holes in the re-forming nuclear envelope requires a fission event to separate the connected INM and ONM (Figure 1a). Whilst localization of ESCRT-III subunits to the telophase nuclei was previously reported [33], recent studies [37^{••},62^{••}] have added mechanistic insights into this process by demonstrating that ESCRT-III plays an essential role in the sealing of the nuclear envelope during late stages of cell division. High-resolution microscopical approaches, including use of Structured-Illumination-Microscopy (SIM) and correlative light and electron microscopy (CLEM), demonstrated the transient recruitment of ESCRT-III components and VPS4 to sites of annular fusion in the reforming nuclear envelope (Figure 3a) [37^{••},62^{••}]. Importantly, these sites of annular fusion comprised membranous stalks connecting INM and ONM and were of similar dimensions to the stalks connecting nascent virions to the plasma membrane or intraluminal vesicles to the endosomal limiting membrane (30–50 nm) [62^{••}]. The transient nature of this recruitment was exposed through live imaging approaches, demonstrating recruitment times of between 1 and 5 min, paralleling well the timings of ESCRT-III recruitment to sites of HIV-1 release at the plasma membrane [63–65]. ESCRT-III assembly on the reforming nuclear envelope occurs through a canonical pathway, with sequential recruitment of CHMP4, CHMP3 and CHMP2 subunits [62^{••}], highlighting its similarities to other ESCRT-III-driven processes. Furthermore, the previously uncharacterized CHMP7 protein was identified as an essential recruiter of CHMP4B during nuclear envelope

reformation, giving at last a role for this protein [37^{••}]. A yeast CHMP7, long thought erroneously to not exist, has recently been implicated in NPC assembly [66] which may further link NPC surveillance and membrane sealing functions at this membrane. CHMP7 is unique amongst ESCRT-III subunits in that it contains an extended N-terminal domain of unknown function — whether this domain specifies a role in nuclear envelope functionality remains to be established.

Importantly, correlative electron tomography analysis demonstrated that ESCRT-III depletion results in the persistence of unsealed holes in the post-mitotic nuclear envelope. As shown in compartmentalization assays, these unsealed nuclear envelopes are functionally leaky, failing to ensure proper nucleocytoplasmic partitioning of a variety of import reporters and ultimately leading to the appearance of DNA damage foci at presumptive unsealed sites, highlighting the important role ESCRT-III has in protecting the genome from damage during mitotic exit [37^{••},62^{••}]. An interaction was also described between the p97 adaptor protein UFD1 and the late-acting ESCRT-III subunit CHMP2A. p97/UFD1 has a previously described role in nuclear envelope sealing [61] and regulates extraction of ubiquitinated Aurora-B from chromatin, allowing chromatin decondensation as the envelope is closed [67,68]. Although CHMP2A is recruited to the forming nuclear envelope through interaction with CHMP4 proteins, UFD1 depletion also impaired recruitment of CHMP2A to this membrane — in this context, UFD1's ability to interact with the auto-inhibitory helix of CHMP2A suggests that this complex may play an additional role in regulating CHMP2A incorporation and assembly of ESCRT-III at the reforming nuclear envelope.

During cytokinesis, the ESCRT-III associated AAA-ATPase Spastin is recruited to the midbody through interaction with CHMP1B and IST1 [35,69,70]. Here, it disassembles midbody microtubules, coordinating the cytoskeletal and membrane remodeling events necessary for abscission. Satisfyingly, during nuclear envelope reformation, Spastin plays a complementary role in disassembling spindle microtubules (which pass through the reforming nuclear envelope), a necessary step before the nuclear envelope is sealed [37^{••}]. Here, the CHMP-like protein IST1 was necessary for recruitment of Spastin and subsequent MT disassembly. Disruption of Spastin results in impaired spindle disassembly and persistence of CHMP4B at the re-forming nuclear envelope, indicating that ESCRT-III function at this membrane cannot be completed until MTs penetrating the reassembling nuclear envelope are removed. These results suggest that ESCRT-III and Spastin work together to coordinate nuclear envelope sealing with disassembly of MTs during mitotic exit and highlight a conservation of the machineries that regulate mitotic membrane remodeling events (Figure 4).

Figure 4



Mitotic co-ordination of membrane and cytoskeletal remodeling by ESCRT-III. A. Cartoon depicting similarities between ESCRT-III dependent nuclear envelope reformation (a) and cytokinesis (b). Enlargement of green-boxed region given in lower cartoon. In both cases, prior to ESCRT-III-dependent membrane resolution, microtubules must be coordinately removed through the action of the AAA-ATPase Spastin, which is recruited to the site of remodeling through interaction with IST1 during nuclear envelope reformation, or IST1/CHMP1B during cytokinesis. In the case of cytokinesis, abscission occurs at a constriction upon the midbody arms and whilst cuts can occur on both sides of the Flemming body, here a single cut is depicted with the asymmetric resorption of the Flemming body by one daughter cell as a midbody remnant.

These recent findings reveal a new localization and function for ESCRT-III in nuclear envelope remodeling at sites of annular fusion. ESCRT-III may play additional roles at the nuclear envelope — for example, viruses and

vesicles containing megaRNPs must transverse this membrane [15,71,72] and whilst current models propose alternate transit mechanisms (for example, involving the inter-membrane AAA-ATPase Torsin [73], or viral

proteins themselves [74]), ESCRT-III may contribute to this transit. Indeed, it is possible that the surveillance role of ESCRT-III at the yeast nuclear envelope may involve coupling of defective nucleoporin extraction with ESCRT-III-dependent membrane repair. Budding profiles and intra-NE vesicles accumulated in both *δvps4* yeast and temperature sensitive *npl4* yeast (a UFD1-interacting co-factor of p97 that has been previously implicated in nuclear envelope sealing) [75], indicating parallels may exist between these two functions of ESCRT-III on the nuclear envelope. These data, together with recently described activities of ESCRT-III in the repair of plasma membrane wounds [22•,23•] points to a role of the ESCRT-III complex in regulating membrane quality — in this case, that of the nuclear envelope. The ESCRT-machinery thus has an important role in the dynamic maintenance of compartmental identity, organelle biogenesis and acts to safeguard the genome from damaging agents. This may be particularly relevant in the context of diseases like laminopathies or cancer, where transient nuclear envelope rupturing and the generation of micronuclei has been reported and is thought to contribute to the disease pathology and development [76–79].

Future perspectives

These are exciting times for ESCRT-biology — originally thought to be a key player in the biogenesis of MVBs, the repertoire of physiological and pathophysiological events that involve ESCRT proteins has expanded greatly. In many ways, this reflects the cell's exquisite ability to conserve and repurpose its machineries to accomplish a variety of functions — in this case, those functions that need a topologically equivalent membrane remodeling. ESCRT-III and VPS4 components are conserved as far-back as *Archaea* (prokaryotic organisms lacking membrane bound organelles and nuclei), suggesting that cell division may be considered an ancestral role for the ESCRT-III machinery that evolved as a mechanism to allow binary fission and has been co-opted throughout evolution to allow acquisition of increasingly complex cellular functions [80]. Still, there are many unknowns; whether a surveillance role for ESCRT-III on the nuclear envelope exists in cells performing open mitosis is currently unknown and how a surveillance role for ESCRT-III is coordinated with ESCRT-III-dependent sealing of the reforming nuclear envelope remains an area of open investigation. ESCRT-III is also a key regulator of an Aurora B-dependent cytokinesis checkpoint that is engaged to delay abscission when mitotic defects (including lagging chromosomes [52,81] and defective NPC assembly [82–84]) are detected and it is possible that a surveillance role for ESCRT-III-dependent extraction of defective nucleoporins is coupled to an inhibition of ESCRT-III-dependent abscission. This abscission checkpoint is Aurora B-dependent, and Aurora B activity has recently been shown to delay closure of holes in the nuclear envelope to allow

lagging chromosomes to be incorporated into forming daughter nuclei [85]. Given the role Aurora B plays in regulating ESCRT-III activity in the context of abscission, it is tempting to speculate that a similar regulation of ESCRT-III may occur during this aspect of nuclear envelope reassembly. Mitotic biology is proving a fruitful area in which to explore ESCRT-function and it is likely that mechanistic insight gleaned from these studies will aid our understanding of this important machinery in non-dividing and pathophysiological states.

Acknowledgments

J.G.C is a Wellcome Trust Research Career Development Fellow (093603/Z/10/Z). We acknowledge the Nikon Imaging Centre at KCL and the NIHR Comprehensive Biomedical Research Centre at Guy's and St Thomas' NHS Foundation Trust for access to core equipment to allow acquisition of images.

References and recommended reading

Papers of particular interest, published within the period of review, have been highlighted as:

- of special interest
- of outstanding interest

1. Katzmann DJ, Babst M, Emr SD: **Ubiquitin-dependent sorting into the multivesicular body pathway requires the function of a conserved endosomal protein sorting complex, ESCRT-I.** *Cell* 2001, **106**:145–155.
2. Babst M, Katzmann DJ, Snyder WB, Wendland B, Emr SD: **Endosome-associated complex, ESCRT-II, recruits transport machinery for protein sorting at the multivesicular body.** *Dev Cell* 2002, **3**:283–289.
3. Babst M, Katzmann DJ, Estepa-Sabal EJ, Meerloo T, Emr SD: **Escrt-III: an endosome-associated heterooligomeric protein complex required for mvb sorting.** *Dev Cell* 2002, **3**:271–282.
4. Wollert T, Wunder C, Lippincott-Schwartz J, Hurley JH: **Membrane scission by the ESCRT-III complex.** *Nature* 2009, **458**:172–177.
5. Wollert T, Hurley JH: **Molecular mechanism of multivesicular body biogenesis by ESCRT complexes.** *Nature* 2010, **464**:864–869.
6. Colombo M, Moita C, van Niel G, Kowal J, Vigneron J, Benaroch P, Manel N, Moita LF, Théry C, Raposo G: **Analysis of ESCRT functions in exosome biogenesis, composition and secretion highlights the heterogeneity of extracellular vesicles.** *J Cell Sci* 2013, **126**:5553–5565.
7. Baietti MF, Zhang Z, Mortier E, Melchior A, Degeest G, Geeraerts A, Ivarsson Y, Depoortere F, Coomans C, Vermeiren E *et al.*: **Syndecan-syntenin-ALIX regulates the biogenesis of exosomes.** *Nat Cell Biol* 2012, **14**:677–685.
8. Samson RY, Bell SD: **Ancient ESCRTs and the evolution of binary fission.** *Trends Microbiol* 2009, **17**:507–513.
9. Garrus JE, Schwedler von UK, Pornillos OW, Morham SG, Zavitz KH, Wang HE, Wettstein DA, Stray KM, Côté M, Rich RL *et al.*: **Tsg101 and the vacuolar protein sorting pathway are essential for HIV-1 budding.** *Cell* 2001, **107**:55–65.
10. Martin-Serrano J, Zang T, Bieniasz PD: **HIV-1 and Ebola virus encode small peptide motifs that recruit Tsg101 to sites of particle assembly to facilitate egress.** *Nat Med* 2001, **7**:1313–1319.

These papers were the first to describe how HIV-1 hijacks the ESCRT-machinery to effect its egress.

11. Demirov DG, Ono A, Orenstein JM, Freed EO: **Overexpression of the N-terminal domain of TSG101 inhibits HIV-1 budding by blocking late domain function.** *Proc Natl Acad Sci U S A* 2002, **99**:955-960.

These papers were the first to describe how HIV-1 hijacks the ESCRT-machinery to effect its egress.

12. VerPlank L, Bouamr F, LaGrassa TJ, Agresta B, Kikonyogo A, Leis J, Carter CA: **Tsg101, a homologue of ubiquitin-conjugating (E2) enzymes, binds the L domain in HIV type 1 Pr55(Gag).** *Proc Natl Acad Sci U S A* 2001, **98**:7724-7729.

These papers were the first to describe how HIV-1 hijacks the ESCRT-machinery to effect its egress.

13. Strack B, Calistri A, Craig S, Popova E, Göttinger HG: **AIP1/ALIX is a binding partner for HIV-1 p6 and EIAV p9 functioning in virus budding.** *Cell* 2003, **114**:689-699.
14. Martin-Serrano J, Yarovoy A, Perez-Caballero D, Bieniasz PD, Yaravoy A: **Divergent retroviral late-budding domains recruit vacuolar protein sorting factors by using alternative adaptor proteins.** *Proc Natl Acad Sci U S A* 2003, **100**:12414-12419.
15. Lee C-P, Liu P-T, Kung H-N, Su M-T, Chua H-H, Chang Y-H, Chang C-W, Tsai C-H, Liu F-T, Chen M-R: **The ESCRT machinery is recruited by the viral BFRF1 protein to the nucleus-associated membrane for the maturation of Epstein-Barr Virus.** *PLoS Pathog* 2012, **8**:e1002904.

16. Pawliczek T, Crump CM: **Herpes simplex virus type 1 production requires a functional ESCRT-III complex but is independent of TSG101 and ALIX expression.** *J Virol* 2009, **83**:11254-11264.

17. Carlton JG, Martin-Serrano J: **Parallels between cytokinesis and retroviral budding: a role for the ESCRT machinery.** *Science* 2007, **316**:1908-1912.

These papers were the first to show that ESCRTs were recruited to the midbody and were necessary for cytokinesis.

18. Morita E, Sandrin V, Chung H-Y, Morham SG, Gygi SP, Rodesch CK, Sundquist WI: **Human ESCRT and ALIX proteins interact with proteins of the midbody and function in cytokinesis.** *EMBO J* 2007, **26**:4215-4227.

These papers were the first to show that ESCRTs were recruited to the midbody and were necessary for cytokinesis.

19. Lee HH, Elia N, Ghirlando R, Lippincott-Schwartz J, Hurley JH: **Midbody targeting of the ESCRT machinery by a noncanonical coiled coil in CEP55.** *Science* 2008, **322**:576-580.

20. Carlton JG, Agromayor M, Martin-Serrano J: **Differential requirements for Alix and ESCRT-III in cytokinesis and HIV-1 release.** *Proc Natl Acad Sci U S A* 2008, **105**:10541-10546.

21. Elia N, Sougrat R, Spurlin TA, Hurley JH, Lippincott-Schwartz J: **Dynamics of endosomal sorting complex required for transport (ESCRT) machinery during cytokinesis and its role in abscission.** *Proc Natl Acad Sci U S A* 2011, **108**:4846-4851.

22. Jimenez AJ, Maiuri P, Lafaurie-Janvore J, Divoux S, Piel M, Perez F: **ESCRT machinery is required for plasma membrane repair.** *Science* 2014, **343**:1247136.

These papers described an unexpected role for ESCRT-III in plasma membrane repair.

23. Scheffer LL, Sreetama SC, Sharma N, Medikayala S, Brown KJ, Defour A, Jaiswal JK: **Mechanism of Ca²⁺-triggered ESCRT assembly and regulation of cell membrane repair.** *Nat Commun* 2014, **5**:5646.

These papers described an unexpected role for ESCRT-III in plasma membrane repair.

24. Filimonenko M, Stuffers S, Raiborg C, Yamamoto A, Malerød L, Fisher EMC, Isaacs A, Brech A, Stenmark H, Simonsen A: **Functional multivesicular bodies are required for autophagic clearance of protein aggregates associated with neurodegenerative disease.** *J Cell Biol* 2007, **179**:485-500.

25. Nara A, Mizushima N, Yamamoto A, Kabeya Y, Ohsumi Y, Yoshimori T: **SKD1 AAA ATPase-dependent endosomal transport is involved in autolysosome formation.** *Cell Struct Funct* 2002, **27**:29-37.

26. Loncle N, Agromayor M, Martin-Serrano J, Williams DW: **An ESCRT module is required for neuron pruning.** *Sci Rep* 2015, **5**:8461.

27. Choudhuri K, Llodrá J, Roth EW, Tsai J, Gordo S, Wucherpennig KW, Kam LC, Stokes DL, Dustin ML: **Polarized release of T-cell-receptor-enriched microvesicles at the immunological synapse.** *Nature* 2014, **507**:118-123.

28. Hurley JH: **ESCRTs are everywhere.** *EMBO J* 2015 <http://dx.doi.org/10.15252/embj.201592484>.

29. McCullough J, Colf LA, Sundquist WI: **Membrane fission reactions of the mammalian ESCRT pathway.** *Annu Rev Biochem* 2013 <http://dx.doi.org/10.1146/annurev-biochem-072909-101058>.

30. Mierzwa B, Gerlich DW: **Cytokinetic abscission: molecular mechanisms and temporal control.** *Dev Cell* 2014, **31**:525-538.

31. Henne WM, Buchkovich NJ, Emr SD: **The ESCRT pathway.** *Dev Cell* 2011, **21**:77-91.

32. Howard TL, Stauffer DR, Degnin CR, Hollenberg SM: **CHMP1 functions as a member of a newly defined family of vesicle trafficking proteins.** *J Cell Sci* 2001, **114**:2395-2404.

33. Stauffer DR, Howard TL, Nyun T, Hollenberg SM: **CHMP1 is a novel nuclear matrix protein affecting chromatin structure and cell-cycle progression.** *J Cell Sci* 2001, **114**:2383-2393.

34. McCullough J, Clippinger AK, Talledge N, Skowrya ML, Saunders MG, Naismith TV, Colf LA, Afonine P, Arthur C, Sundquist WI et al.: **Structure and membrane remodeling activity of ESCRT-III helical polymers.** *Science* 2015 <http://dx.doi.org/10.1126/science.1258305>.

This paper was the first to describe the structure of an ESCRT-III subunit in the open conformation.

35. Agromayor M, Carlton JG, Phelan JP, Matthews DR, Carlin LM, Ameer-Beg S, Bowers K, Martin-Serrano J: **Essential role of H1ST1 in cytokinesis.** *Mol Biol Cell* 2009, **20**:1374-1387.

36. Bajorek M, Morita E, Skalicky JJ, Morham SG, Babst M, Sundquist WI: **Biochemical analyses of human IST1 and its function in cytokinesis.** *Mol Biol Cell* 2009, **20**:1360-1373.

37. Vietri M, Schink KO, Campsteijn C, Wegner CS, Schultz SW, Christ L, Thoresen SB, Brech A, Raiborg C, Stenmark H: **Spastin and ESCRT-III coordinate mitotic spindle disassembly and nuclear envelope sealing.** *Nature* 2015, **522**:231-235.

These papers described an essential role for the ESCRT-III complex in resealing the reforming nuclear envelope during mitotic exit.

38. Allison R, Lumb JH, Fassier C, Connell JW, Martin Ten D, Seaman MNJ, Hazan J, Reid E: **An ESCRT-spastin interaction promotes fission of recycling tubules from the endosome.** *J Cell Biol* 2013, **202**:527-543.

39. Adell MAY, Teis D: **Assembly and disassembly of the ESCRT-III membrane scission complex.** *FEBS Lett* 2011, **585**:3191-3196.

40. Fabrikant G, Lata S, Riches JD, Briggs JAG, Weissenhorn W, Kozlov MM: **Computational model of membrane fission catalyzed by ESCRT-III.** *PLoS Comput Biol* 2009, **5**:e1000575.

41. Hurley JH, Hanson PI: **Membrane budding and scission by the ESCRT machinery: it's all in the neck.** *Nat Rev Mol Cell Biol* 2010, **11**:556-566.

42. Shen Q-T, Schuh AL, Zheng Y, Quinney K, Wang L, Hanna M, Mitchell JC, Otegui MS, Ahlquist P, Cui Q et al.: **Structural analysis and modeling reveals new mechanisms governing ESCRT-III spiral filament assembly.** *J Cell Biol* 2014, **206**:763-777.

These papers describe a new way of considering how the spiral ESCRT-III polymer may remodel membranes.

43. Chiaruttini N, Redondo-Morata L, Colom A, Humbert F, Lenz M, Scheuring S, Roux A: **Relaxation of loaded ESCRT-III spiral springs drives membrane deformation.** *Cell* 2015, **163**:866-879.

These papers describe a new way of considering how the spiral ESCRT-III polymer may remodel membranes.

44. Guizetti J, Schermelleh L, Mäntler J, Maar S, Poser I, Leonhardt H, Müller-Reichert T, Gerlich DW: **Cortical constriction during**

- abscission involves helices of ESCRT-III-dependent filaments.** *Science* 2011, **331**:1616-1620.
45. Reid E, Connell J, Edwards TL, Duley S, Brown SE, Sanderson CM: **The hereditary spastic paraplegia protein spastin interacts with the ESCRT-III complex-associated endosomal protein CHMP1B.** *Hum Mol Genet* 2005, **14**:19-38.
 46. Kabachinski G, Schwartz TU: **The nuclear pore complex — structure and function at a glance.** *J Cell Sci* 2015, **128**:423-429.
 47. Webster BM, Colombi P, Jäger J, Lusk CP: **Surveillance of nuclear pore complex assembly by ESCRT-III/Vps4.** *Cell* 2014, **159**:388-401.
- This paper was the first to describe a role for ESCRT-III on the nuclear envelope — in surveillance and extraction of defective nucleoporins.
48. Yewdell WT, Colombi P, Makhnevych T, Lusk CP: **Lumenal interactions in nuclear pore complex assembly and stability.** *Mol Biol Cell* 2011, **22**:1375-1388.
 49. Foresti O, Rodriguez-Vaello V, Funaya C, Carvalho P: **Quality control of inner nuclear membrane proteins by the Asi complex.** *Science* 2014, **346**:751-755.
 50. Khmelinskii A, Blaszcak E, Pantazopoulou M, Fischer B, Omrus DJ, Le Dez G, Brossard A, Gunnarsson A, Barry JD, Meurer M et al.: **Protein quality control at the inner nuclear membrane.** *Nature* 2014, **516**:410-413.
 51. Webster BM, Lusk CP: **ESCRTs breach the nuclear border.** *Nucleus* 2015, **6**:197-202.
 52. Carlton JG, Caballe A, Agromayor M, Kloc M, Martin-Serrano J: **ESCRT-III governs the Aurora B-mediated abscission checkpoint through CHMP4C.** *Science* 2012, **336**:220-225.
 53. Güttinger S, Laurell E, Kutay U: **Orchestrating nuclear envelope disassembly and reassembly during mitosis.** *Nat Rev Mol Cell Biol* 2009, **10**:178-191.
 54. Schooley A, Vollmer B, Antonin W: **Building a nuclear envelope at the end of mitosis: coordinating membrane reorganization, nuclear pore complex assembly, and chromatin de-condensation.** *Chromosoma* 2012, **121**:539-554.
 55. Burke B: **The nuclear envelope: filling in gaps.** *Nat Cell Biol* 2001, **3**:E273-E274.
 56. Burke B, Ellenberg J: **Remodelling the walls of the nucleus.** *Nat Rev Mol Cell Biol* 2002, **3**:487-497.
 57. Lu L, Ladinsky MS, Kirchhausen T: **Cisternal organization of the endoplasmic reticulum during mitosis.** *Mol Biol Cell* 2009, **20**:3471-3480.
 58. Lu L, Ladinsky MS, Kirchhausen T: **Formation of the postmitotic nuclear envelope from extended ER cisternae precedes nuclear pore assembly.** *J Cell Biol* 2011, **194**:425-440.
 59. Anderson DJ, Hetzer MW: **Nuclear envelope formation by chromatin-mediated reorganization of the endoplasmic reticulum.** *Nat Cell Biol* 2007, **9**:1160-1166.
 60. Puhka M, Joensuu M, Vihinen H, Belevich I, Jokitalo E: **Progressive sheet-to-tubule transformation is a general mechanism for endoplasmic reticulum partitioning in dividing mammalian cells.** *Mol Biol Cell* 2012, **23**:2424-2432.
 61. Hetzer M, Meyer HH, Walther TC, Bilbao-Cortes D, Warren G, Mattaj JW: **Distinct AAA-ATPase p97 complexes function in discrete steps of nuclear assembly.** *Nat Cell Biol* 2001, **3**:1086-1091.
- These papers described an unexpected role for ESCRT-III in plasma membrane repair.
62. Olmos Y, Hodgson L, Mantell J, Verkade P, Carlton JG: **ESCRT-III controls nuclear envelope reformation.** *Nature* 2015, **522**:236-239.
- These papers described an essential role for the ESCRT-III complex in resealing the reforming nuclear envelope during mitotic exit.
63. Jouvenet N, Zhadina M, Bieniasz PD, Simon SM: **Dynamics of ESCRT protein recruitment during retroviral assembly.** *Nat Cell Biol* 2011, **13**:394-401.
 64. Baumgärtel V, Ivanchenko S, Dupont A, Sergeev M, Wiseman PW, Kräusslich H-G, Bräuchle C, Müller B, Lamb DC: **Live-cell visualization of dynamics of HIV budding site interactions with an ESCRT component.** *Nat Cell Biol* 2011, **13**:469-474.
 65. Bleck M, Itano MS, Johnson DS, Thomas VK, North AJ, Bieniasz PD, Simon SM: **Temporal and spatial organization of ESCRT protein recruitment during HIV-1 budding.** *Proc Natl Acad Sci U S A* 2014, **111**:12211-12216.
 66. Bauer I, Brune T, Preiss R, Kölling R: **Evidence for a nonendosomal function of the *Saccharomyces cerevisiae* ESCRT-III-like protein Chm7.** *Genetics* 2015, **201**:1439-1452.
 67. Ramadan K, Bruderer R, Spiga FM, Popp O, Baur T, Gotta M, Meyer HH: **Cdc48/p97 promotes reformation of the nucleus by extracting the kinase Aurora B from chromatin.** *Nature* 2007, **450**:1258-1262.
 68. Dobrynin G, Popp O, Romer T, Bremer S, Schmitz MHA, Gerlich DW, Meyer H: **Cdc48/p97-Ufd1-Npl4 antagonizes Aurora B during chromosome segregation in HeLa cells.** *J Cell Sci* 2011, **124**:1571-1580.
 69. Connell JW, Lindon C, Luzio JP, Reid E: **Spastin couples microtubule severing to membrane traffic in completion of cytokinesis and secretion.** *Traffic* 2009, **10**:42-56.
 70. Yang D, Rismanchi N, Renvoisé B, Lippincott-Schwartz J, Blackstone C, Hurley JH: **Structural basis for midbody targeting of spastin by the ESCRT-III protein CHMP1B.** *Nat Struct Mol Biol* 2008, **15**:1278-1286.
 71. Speese SD, Ashley J, Jokhi V, Nunnari J, Barria R, Li Y, Ataman B, Koon A, Chang Y-T, Li Q et al.: **Nuclear envelope budding enables large ribonucleoprotein particle export during synaptic Wnt signaling.** *Cell* 2012, **149**:832-846.
 72. Rose A, Schlieker C: **Alternative nuclear transport for cellular protein quality control.** *Trends Cell Biol* 2012, **22**:509-514.
 73. Jokhi V, Ashley J, Nunnari J, Noma A, Ito N, Wakabayashi-Ito N, Moore MJ, Budnik V: **Torsin mediates primary envelopment of large ribonucleoprotein granules at the nuclear envelope.** *Cell Rep* 2013, **3**:988-995.
 74. Bigalke JM, Heuser T, Nicastro D, Heldwein EE: **Membrane deformation and scission by the HSV-1 nuclear egress complex.** *Nat Commun* 2014, **5**:4131.
 75. DeHoratius C, Silver PA: **Nuclear transport defects and nuclear envelope alterations are associated with mutation of the *Saccharomyces cerevisiae* NPL4 gene.** *Mol Biol Cell* 1996, **7**:1835-1855.
 76. De Vos WH, Houben F, Kamps M, Malhas A, Verheyen F, Cox J, Manders EMM, Verstraeten VLRM, van Steensel MAM, Marcelis CLM et al.: **Repetitive disruptions of the nuclear envelope invoke temporary loss of cellular compartmentalization in laminopathies.** *Hum Mol Genet* 2011, **20**:4175-4186.
 77. Gupta P, Bilinska ZT, Sylvius N, Boudreau E, Veinot JP, Labib S, Bolongo PM, Hamza A, Jackson T, Ploski R et al.: **Genetic and ultrastructural studies in dilated cardiomyopathy patients: a large deletion in the lamin A/C gene is associated with cardiomyocyte nuclear envelope disruption.** *Basic Res Cardiol* 2010, **105**:365-377.
 78. Vargas JD, Hatch EM, Anderson DJ, Hetzer MW: **Transient nuclear envelope rupturing during interphase in human cancer cells.** *Nucleus* 2012, **3**:88-100.
 79. Hatch EM, Fischer AH, Deerinck TJ, Hetzer MW: **Catastrophic nuclear envelope collapse in cancer cell micronuclei.** *Cell* 2013, **154**:47-60.
 80. Baum DA, Baum B: **An inside-out origin for the eukaryotic cell.** *BMC Biol* 2014, **12**:76.
 81. Steigemann P, Wurzenberger C, Schmitz MH, Held M, Guizetti J, Maar S, Gerlich DW: **Aurora B-mediated abscission checkpoint protects against tetraploidization [Internet].** *Cell* 2009, **136**:473-484.
 82. Mackay DR, Makise M, Ullman KS: **Defects in nuclear pore assembly lead to activation of an Aurora B-mediated abscission checkpoint.** *J Cell Biol* 2010, **191**:923-931.

83. Mackay DR, Elgort SW, Ullman KS: **The nucleoporin Nup153 has separable roles in both early mitotic progression and the resolution of mitosis.** *Mol Biol Cell* 2009, **20**: 1652-1660.
84. Caballe A, Wenzel DM, Agromayor M, Alam SL, Skalicky JJ, Kloc M, Carlton JG, Labrador L, Sundquist WI, Martin-Serrano J: **ULK3 regulates cytokinetic abscission by phosphorylating ESCRT-III proteins.** *Elife* 2015, **4**:e06547.
85. Karg T, Warecki B, Sullivan W: **Aurora B-mediated localized delays in nuclear envelope formation facilitate inclusion of late-segregating chromosome fragments.** *Mol Biol Cell* 2015, **26**:2227-2241.